




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 9/02, 47/34, 47/36</b>		A1	(11) International Publication Number: <b>WO 97/30693</b>
			(43) International Publication Date: 28 August 1997 (28.08.97)
(21) International Application Number: <b>PCT/KR97/00032</b>		(81) Designated States: CA, CN, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 25 February 1997 (25.02.97)			
(30) Priority Data: 1996/4566 26 February 1996 (26.02.96) KR		Published With international search report.	
(71) Applicant: DONG WHA PHARMACEUTICAL INDUSTRIAL CO., LTD. [KR/KR]; 5, Soonwha-dong, Joong-ku, Seoul 100-130 (KR).			
(72) Inventors: YOON, Sung, June; 1420-11, Sinlim 5-dong, Kwanac-ku, Seoul 151-015 (KR). RYU, Jei, Man; 136-103, Kwanac Apartment, Buheung-dong, Dongan-ku, Anyang, Kyunggi-do 431-054 (KR). CHOI, Han, Gon; 1-1001, Daelim Apartment, Sadang 3-dong, Dongjak-ku, Seoul 156-093 (KR). JUNG, Jae, Hee; 1326-603, Mogdong Apartment, Sinjung 6-dong, Yangcheon-ku, Seoul 158-076 (KR). SUNG, Yong, Kiel; 317-1601, Woosung Apartment, Jungja-dong, Bundang-ku, Sungnam, Kyunggi-do 463-010 (KR). YOO, Jong, Ho; 2-809, Samho Apartment, Bisan 3-dong, Dongan-ku, Anyang, Kyunggi-do 431-053 (KR).			
(74) Agent: SUH, Jong, Wan; New-Seoul Building, 3rd floor, 828-8, Yeoksam-dong, Kangnam-ku, Seoul 135-080 (KR).			

(54) Title: SUPPOSITORY COMPOSITION OF THE DRUG WHICH UNDERGO THE HEPATIC FIRST-PASS EFFECT

## (57) Abstract

This invention relates to a suppository composition containing a drug which undergoes the hepatic first-pass effect, poloxamer and hydrophilic natural polymers. The suppository composition of this invention is characterized in that: has the gelation temperature of 30 to 36 °C, and is a liquid form at room temperature, and readily becomes a gel at body temperature after rectal administration; has the remarkable gel strength, and is not leaked out the anus; has the remarkable bioadhesive force, and doesn't climb up to the end of the colon, therefore ensures better bioavailability of the drug.



- - intravenous injection group of propranolol solution
- - oral administration group of propranolol solution
- Δ - rectal administration group of conventional suppository composition 1
- ▼ - rectal administration group of the comparative suppository composition
- - rectal administration group of the suppository composition of Example 2

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## THE TITLE OF INVENTION

Suppository composition of the drug  
which undergo the hepatic first-pass effect

5

### TECHNICAL FIELD

This invention relates to the suppository composition of the drug which undergo the hepatic first-pass effect and more particularly, a suppository composition of the drug which undergo the hepatic first-pass effect, being characterized in that said composition;

10

- has the gelation temperature of 30 to 36 °C, and is a liquid form at room temperature, and readily becomes a gel at body temperature after rectal administration;

- has the remarkable gel strength, and is not leaked out from the anus;

15

- has the remarkable bioadhesive force, and doesn't climb upto the end of the colon, therefore ensures better bioavailability of the drug.

### BACKGROUND ART

The oral drug administration is the commonest method designed to apply the drugs into the human body. However, some drugs are absorbed from the gastrointestinal tract and eliminated by the hepatic first-pass effect, thus giving lower bioavailability and significant reduction of therapeutic efficacy thereto. The drugs which undergo the hepatic first-pass effect include propranolol, alprenol, metoprolol, isoproterenol, oxprenolol, pindolol, testosterone, methyltestosterone, epinephrine, imipramine, desmethylinipramine, nortryptiline, oxyphenbutazone, tryptophan, serotonin, pheniprazine, morphine, lidocain, propoxyphene, salicylamide, hexobarbital, phenyltoin, quaternary ammonium salt, pentazocine, diltiazem, nifedipine, nitrandipine, and cyclosporin A. In an effort to enhance their bioavailability, these drugs in an injectable form are being given but it has recognized several disadvantages in that a)

20  
25  
30

administration is inconvenient, b) most of patients dislike an injection, and c) drug's direct infusion into the blood vessel is very dangerous.

In order to overcome these shortcomings as aforementioned, there have been several approaches to change the administration route of drugs which undergo the hepatic first-pass effect, into other alternative routes via nose, skin, rectum or etc. However, said administration routes cannot completely meet the desired objectives in that;

- Nasal administration of propranolol is deemed inconvenient for the handling and administration of dosage form(Hussain et al., *J. Pharm. Sci.*, 69, 1411-1413, 1980);

- The possible skin-permeation tests of drug using multilaminate adhesive device or iontophoresis show that the amount of drug permeated through skin is so small that the transdermal administration of drug cannot achieve the target dose(Corbo et al., *Pharm. Res.*, 6(9), 753-758, 1989; Singh et al., *J. Control. Rel.*, 18, 165-170, 1992);

- Rectal administration when dosage is given induces disgusting feeling and discomforts and after administration the dosage climbs upto the end of the colon by peristalsis, therefore the drug is absorbed there and undergo the hepatic first-pass effect. Thus said administration route has failed to get over the aforementioned defects completely.

In case of testosterone, there have been some approaches of the buccal and the transdermal administration but failed due to insufficient amount of drugs permeated through mouth mucosa and skin.

The inventor et al. have noticed the rectal administration of drugs which undergo the hepatic first-pass effect and in order to get over the aforementioned shortcomings associated with said rectal administration, poloxamer as suppository base is under careful consideration since such base is in a liquid phase at low temperature but when temperature goes up, it becomes in a gel phase.

In recent years, considerable interest has focused on some approaches to apply the poloxamer into the human body. For example the

U.S. Pat. No. 4,188,373 discloses the adjustment of gelation temperature depending on varied concentrations of poloxamer and the Canadian Pat. No. 1,072,413 also discloses varied gelatin temperatures of bases through mixing with Tetronic® and Tergitol®, poloxamer derivatives.

5        Several U.S. Pat. Nos. 4,478,822, 4,474,751, 4,474,752 and 4,474,753 also disclose that using poloxamer as a base, various kinds of additive are employed so as to adjust the ion strength and pH as well as to apply it into skin, eye and the body cavity such as rectum and urinary tract.

10        However, the conventional gel composition using said poloxamer has some disadvantages in that when rectal administration is given, due to undesirable conditions of gelation temperature, gel strength and bioadhesive force, the dosage may be leaked out from the anus, or climb upto the end of the colon. Thus the drug of that dosage form may not be  
15        absorbed, or that absorbed at colon undergo the hepatic first-pass metabolism.

      The Europe Patent No. 0 551 626 A1 discloses that with pH adjustment and addition of carbomer, said poloxamer with increased gelation temperature and viscosity is applied to skin, eye, rectum and  
20        urinary tract. Nevertheless, this invention has proven insufficient in its application to the suppository composition since it does not consider the bioadhesive force and dissolution.

## DISCLOSURE OF THE INVENTION

25        In order to overcome the aforementioned shortcomings, therefore, the inventor et al. have endeavored to develop a suppository composition of the drug which undergo the hepatic first-pass effect with better bioavailability.

      An object of this invention is to provide a suppository composition  
30        of the drug which undergo the hepatic first-pass effect, being characterized in that;

- Any disgusting feeling or discomforts does not occur, when a drug is administered;
- Administration is easy and after administration a composition is not leaked out from the anus or does not climb upto the end of the colon, thus ensuring better bioavailability.

The suppository composition of this invention comprises 0.1 ~ 10 weight part of a drug which undergoes the hepatic first-pass effect, 25 ~ 40 weight part of poloxamer and 0.1 ~ 1 weight part of hydrophilic natural polymers to the total blending ratio.

The drug, contained in the composition of this invention, which undergo the hepatic first-effect, may be selected from the following materials: propranolol, alprenol, metoprolol, isoproterenol, oxprenolol, pindolol, testosterone, methyltestosterone, epinephrine, imipramine, desmethyylimipramine, nortryptiline, oxyphenbutazone, tryptophan, serotonin, pheniprazine, morphine, lidocain, propoxyphene, salicylamide, hexobarbital, phenyltoin, quarternary ammonium salt, pentazocine, diltiazem, nifedipine, nitrandipine, cyclosporin A and etc.. In case of the amount of this drug, 0.1 ~ 10 weight part is contained to the total suppository composition; If less than 0.1 weight part is contained, relatively enlarged volume of said composition at single dose makes it difficult to perform the rectal administration but in case of exceeding 10 weight part, the drug itself reduces the gel strength and bioadhesive force so that the adjustment of that properties become difficult.

From the suppository composition of this invention, one or more poloxamers may be selected from the following, i.e., solid-phase type (F-127, F-108, F-98, F-88, F-68 and etc.), liquid-phase type (L-44, L-62, L-64 and etc.) and paste type (P-85, P-81, P-123 and etc.). It is preferred to contain 25 ~ 40 weight part of poloxamer to the suppository composition; if less than 25 weight part is contained, the gel strength and bioadhesive force are weak and in case of exceeding 40 weight part, higher degree of

viscosity makes it difficult to manufacture the desired product.

These poloxamers adjust the gelation temperature of the suppository composition to 30~36 °C; thus, said poloxamers are in liquid phase at room temperature and in gel state within the body.

5       The hydrophilic natural polymers contained in the suppository composition of this invention may be used by selecting chitosan and sodium alginate independently or in a mixing form.

      About 0.1~1 weight part of hydrophilic natural polymers is contained to the suppository composition; if less than 0.1 weight part is  
10       contained, the gel strength and bioadhesive force cannot be adjusted and sustained drug release is unavailable. Meantime, in case of exceeding 1 weight part, higher degree of viscosity makes it difficult to manufacture the desired product.

      Since these hydrophilic natural polymers have hydrophilic groups  
15       such as amine group (NH<sub>2</sub>) or hydroxyl group (OH) at the end of the molecular structure which may be reacted with hydroxyl group (OH) of poloxamer by hydrogen bond, more strong three-dimensional networking structure of poloxamer may be formed. In addition to that, these polymers can form the strong hydrogen bond with oligosaccharide groups  
20       of rectal mucosa. Therefore, very small amount of those can play a role to reinforce the gel strength and bioadhesive force. Further these hydrophilic natural polymers release the drug slowly by their matrix formation at a constant concentration.

      In addition to the drugs which undergo the hepatic first-pass effect,  
25       poloxamer and hydrophilic natural polymers, the suppository composition of this invention may also include the following additives which may be commonly applied to the conventional dosage form of rectal administration: preservatives (e.g., sodium benzoate, potassium sorbate, paraben derivatives and etc.), pH modulator (e.g., hydrochloric acid, citric acid, sodium hydroxide and etc.), stabilizers (e.g., methionine,  
30       etc.) and etc..

The suppository composition according to this invention may be prepared by dissolving these compositions in an appropriate amount of water.

The suppository composition of this invention is characterized in that;

- has the gelation temperature of 30 to 36 °C, and is a liquid form at room temperature, and readily becomes a gel at body temperature after rectal administration;

- has the remarkable gel strength, and is not leaked out the anus;

- has the remarkable bioadhesive force, and doesn't climb upto the end of the colon, therefore ensures better bioavailability of the drug. Therefore, the suppository composition of this invention may improve a poor bioavailability that the conventional suppository composition faces, i.e., after rectal administration a composition climb upto the end of the colon by peristalsis of the large intestine, and the drug is absorbed there and undergo the hepatic first-pass effect.

With simple manufacturing process, the suppository composition may be easily manufactured with cost-saving effects.

## **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is a graph showing the gelation temperature, when polymers are added.

Fig. 2 is a graph showing the gel strength, when polymers are added.

Fig. 3 is a graph showing the bioadhesive force, when polymers are added.

Fig. 4 is a graph showing the dissolution-controlling capacity, when polymers are added.

Fig. 5 is a graph showing the results of dissolution tests related to the suppository composition of Example 1 ~ 4.

Fig. 6 shows the plasma concentration of propranolol, when



suppository composition of Example 2, comparative suppository composition and conventional suppository compositions are administered via rectum and propranolol solution are administered via injection and mouth, respectively.

5

### **BEST MODE FOR CARRING OUT THE INVENTION**

This invention is explained in more detail by the following examples, but the claims are not limited to these examples.

#### 10 **EXAMPLE 1-4**

The blending ratio involving the suppository composition according to Example 1-4 is shown in the following table 1. In the blending ratio as shown hereunder, poloxamer and hydrophilic natural polymers were dissolved in water and then, drugs and other components  
15 were successively added to the mixture for dissolving completely. Then, water was added to be a total of 100g in the weight of this mixture and the suppository composition was finally prepared.

Table 1. Blending ratio for the suppository composition according to Example 1-4

composition		example 1	example 2	example 3	example 4
poloxamer	F-127	15	15		20
	F-108			12	
	F-88			18	20
	F-68	19	15		
hydrophilic natural polymer	sodium alginate	02	06		
	chitosan			05	08
drug	propranolol	0.4	2.0		
	testosterone			0.1	
	epinephrine				8.0
additives	methyl parahydroxybenzoate	0.06		0.06	
	propyl parahydroxybenzoate	0.03		0.03	
	sodium benzoate				0.1
	citric acid				0.01
water		appropriate	appropriate	appropriate	appropriate
total (g)		100.0	100.0	100.0	100.0

#### COMPARATIVE EXAMPLE 1-4

5 In the same manner as described in the above compositions of Example 1-4, each composition without addition of hydrophilic natural polymers was prepared.

### EXPERIMENTAL EXAMPLE 1: Selection of polymers adaptable to a suppository composition

For the selection of some polymers adaptable to suppository composition of propranolol, each of the following materials by 1 weight part such as polyvinylpyrrolidone (PVP), hydroxypropylmethylcellulose (HPMC), hydroxypropylcellulose (HPC), carbopol, polycarbophil, sodium alginate and chitosan was added to the mixing solution of poloxamer [F-127/F-68(15/15 weight part)], thus manufacturing the suppository compositions. The gelation temperature, gel strength and bioadhesive force related to each composition were measured. Further 2 weight part of each propranolol were added to the suppository composition, so prepared, so as to measure the dissolution rate of propranolol.

The testing criteria on the gelation temperature, gel strength, bioadhesive force and dissolution rate were as follows:

Gelation temperature: 10g of sample of the suppository composition was charged to a 20ml container, together with magnetic bar and installed to water bath at 4 °C. With a digital thermometer inserted into the sample so as not to contact with the magnetic bar, the sample was stirred at a constant rate and while increasing its temperature at a rate of 1 °C/min, the gelation temperature was determined when the magnetic bar was completely stopped.

Gel strength: 50g of the suppository composition was charged to a 100ml-mass cylinder and equilibrated in water bath at 36.5 °C for 30 mins. With a gel strength device placed on a mass cylinder, the gel strength was determined by time (second) when the device went down to 5cm.

Bioadhesive force: Two sections of tissue cut from the rectal mucosa of rabbit were attached to two vials of a bioadhesive force device and between them, 0.05g of the suppository composition was added. Then, with counterpoises piled up successively, the counterpoise weight

when said vials fall was calculated as a force extended per unit area.

Dissolution rate: 1g of the suppository composition was charged to a semi-permeable membrane and with both sides fastened with threads, the dissolution test was performed in phosphate buffer solution of pH 6.8 at 100 rpm using the paddle method. The small amount of medium was sampled at one hour intervals for analysis thereof.

These results are shown in the following table 1 and Fig. 1-4.

Fig. 1 is a graph showing the effect of the kinds of polymers on the gelation temperature. The gelation temperature of suppository composition was somewhat affected by the polymers irrespective of their kinds.

Fig. 2 is a graph showing the effect of the kinds of polymers on the gel strength and Fig. 3 is a graph showing the effect of the kinds of polymers on the bioadhesive force.

Compared with the suppository composition containing polyvinylpyrrolidone, hydroxypropylmethylcellulose, hydroxypropyl cellulose, carbopol or polycarbophil, the sodium alginate- or chitosan-containing suppository composition showed remarkably high gel strength and bioadhesive force.

Fig. 4 is a graph comparing the dissolution rate of propranolol from the suppository composition containing a certain scope of 0.2~0.8% concentration of sodium alginate and polycarbophil, respectively as hydrophilic polymers. While the suppository composition containing 0.2% sodium alginate showed higher dissolution rate than the polycarbophil-containing composition in same concentration, the composition containing 0.8% sodium alginate showed lower dissolution rate than the polycarbophil-containing composition in same concentration. This revealed that the composition containing a certain scope of 0.2~0.8% of sodium alginate as polymers had a wider range of propranolol dissolution than polycarbophil-containing composition, thus showing superior dissolution-controlling capacity of the former as a hydrophilic

polymer.

In the same manner as described in the above, some tests ascertained that carbopol-containing composition has similar dissolution range of polycarbophil-containing composition and chitosan-containing composition has similar dissolution range of sodium alginate-containing composition.

Since it was judged that sodium alginate and chitosan is superior to other polymers in terms of the gel strength, bioadhesive force and dissolution-controlling capacity. Therefore, the inventor et al. have selected sodium alginate and chitosan as polymers adoptable to the suppository composition of this invention.

EXPERIMENTAL EXAMPLE 2 : Measurement of gelation temperature, gel strength, leakage of the composition from the anus

The gelation temperature, gel strength and bioadhesive force of Comparative example 1-4 and Example 1-4 were measured in a same method as described in Experimental example 1. The animal experiments (assessment on leakage of the composition from the anus) were performed as follows:

- 1g of the suppository composition was inserted into the anus of rabbits in 5cm depth using a stomach sonde needle for rat and with rabbits placed at 45° slope obliquely, observations for 30 mins were made and the judgment was acceptable, when the drug suppository composition was not leaked out from the anus. Its results were shown in the following table

2.

Table 2. Measurement of gelation temperature, gel strength, leakage of the composition from the anus and bioadhesive force related to compositions of Comparative example 1-4 and of Example 1-4

experimental group	gelation temperature (°C)	gel strength (sec)	bioadhesive force (dyne/cm <sup>2</sup> × 10 <sup>2</sup> )	leakage of the composition from the anus
example 1	33.4	16.0	108.0	No
comparative example 1	35.0	8.3	25.7	Yes
example 2	32.2	51.7	152.3	No
comparative example 2	36.9	8.6	12.3	Yes
example 3	31.0	32.4	158.7	No
comparative example 3	34.7	4.3	15.6	Yes
example 4	33.1	19.2	62.6	No
comparative example 4	36.3	9.5	14.4	Yes

The above results showed that the suppository composition of Example 1-4 had lower gelation temperature, better gel strength/bioadhesive force and less leakage of the composition from the anus, than the composition of comparative example 1-4 without hydrophilic natural polymers

#### EXPERIMENTAL EXAMPLE 3: Dissolution test

Each dissolution rate related to 2g of the suppository composition of Example 1-2 and 1g of the suppository composition of Example 3-4 was measured in a same manner as described in Experimental example 1.

The experimental results were shown in Fig. 5 and the suppository composition of Example 1-4 has proven sufficient as a preparation.

#### EXPERIMENTAL EXAMPLE 4: Damage test of mucosal membrane

In order to assess each damage of the mucosal membrane induced by the suppository composition of Example 2, poloxamer, PEG 4000, Witepsol, comparative suppository composition [F-127/F-68/propranolol(15/15/2 weight part)], conventional suppository composition I [PEG 4000/propranolol (98/2 weight part)] and conventional suppository composition II [Witepsol/propranolol (98/2 weight part)], each group of 3 Sprague-Dawley rats, being fasted for 24-36 hours, was used for the following experiment.

Each suppository composition (0.1g) was inserted into the anus of rat in 5cm depth using a stomach sonde needle for rat. After 4 hrs, the rectum was harvested, cut by a knife and cleansed with physiological saline solution. The rectum was fixed with formaldehyde solution neutralized with 10% carbonate, followed by paraffin and stained with hematoxylin-eosin for microscopical observation.

The damage degree of the rectal mucosa was measured by the method of Reid et al. (Reid, A.S., et al., *Int. J. Pharm.*, 40, 181-185 (1987)), in which the following standards were applied:

Normal: The epithelium at interglandular site was normal.

Type I : Part of epithelium at interglandular site was torn apart or is tearing down.

Type II : The height of epithelium becomes generally low.

Type III : The epitheliums were completely torn apart, exposing the mucosae.

As the type passed into I, II and III, they showed the damage severity of the rectal mucosa. The value was calculated by percent (%). Its results were shown in table 3.

Table 3. Damage rate of rectal mucosa per type (Unit: %)

experimental group	normal	type I	type II	type III	total
normal mucosa	85.7±4.7	4.2±2.9	8.0±3.1	2.1±1.3	100
PEG 4000	60.4±4.5	7.0±2.7	20.6±7.8	12.0±3.3	100
Witepsol	16.0±6.6	8.0±2.6	20.4±4.6	55.6±12.3	100
F-127/F-68 (15/15%)	79.4±7.8	4.0±2.4	9.9±4.0	6.7±3.2	100
conventional suppository composition I	67.6±9.0	8.9±4.4	18.5±5.8	5.0±4.6	100
conventional suppository composition II	9.2±5.6	7.7±5.1	26.3±5.2	56.8±6.6	100
comparative suppository composition	44.9±9.1	7.2±2.6	27.6±8.1	20.3±11.6	100
example 2	72.3±7.7	5.0±1.6	15.9±7.9	6.8±4.9	100

From the above results, poloxamer as bases of the suppository composition of this invention showed more significant reduction in the damage rates of rectal mucosa than PEG 4000 and Witepsol as bases of conventional suppository and the suppository composition showed also less damage rates of rectal mucosa than conventional suppository compositions.

#### 10 EXPERIMENTAL EXAMPLE 5: Location of composition after rectal administration

After rectal administration of the suppository composition of Example 2, comparative suppository composition and conventional suppository composition I, each location was measured as follows:

15 Each suppository composition(0.2g) containing 0.1% Blue No. 1 Lake coloring agent was inserted into the anus of rat in 5cm depth using a stomach sonde needle for rat. The rectums of rats were harvested at intervals of 5 min, 2 hrs and 4 hrs so as to ascertain the location of the



suppository composition. Its results were shown in the following table 4.

Table 4. Location of each composition after rectal administration

experimental group	5 min	2 hrs	4 hrs
conventional suppository composition I	4-5 cm	4-5 cm	7-8 cm
comparative suppository composition	4-5 cm	7-8 cm	7-8 cm
example 2	4-5 cm	4-5 cm	4-5 cm

5        The suppository composition of Example 2 was found at the place 4~5cm away from the anus even after 4 hrs, thus reflecting that said composition did not climb upto the large intestine, while the comparative suppository composition and conventional suppository composition I were found at the place 7~8cm away from the anus at the intervals of 2  
10       hrs and 4 hrs, respectively, thus showing that said composition did not climb upto the large intestine.

#### EXPERIMENTAL EXAMPLE 6: Pharmacokinetics in suppository composition

15        Male rabbits, fasted for 24-36 hours, were used as experimental animals. In order to investigate the pharmacokinetics, the experimental animals were divided into the following 5 groups with each group containing 4 rabbits: i) intravenous injection group of propranolol solution(propranolol 20mg/water for injection 20ml), ii) oral  
20       administration group of propranolol solution(propranolol 20mg/water 20ml), iii) rectal administration group of conventional suppository composition I, iv) rectal administration group of the comparative suppository composition, and v) rectal administration group of the suppository composition of Example 2.

Rats were anesthetized with urethane and fixed them on a fixing stand. Then polyethylene tube was inserted into the right femoral artery and the suppository composition with a dose of propranolol 2mg/kg was given intravenously, orally or rectally. 0.5ml of blood sample from the right femoral artery was collected at certain intervals, centrifuged at 3000 rpm for 30 mins and harvested 0.2ml of plasma. 0.2ml of internal standard-acetonitrile(200mcg/ml) was added to the plasma and centrifuged at 3000 rpm for 10 mins to precipitate protein. The resulting solution was analyzed by HPLC. Its results were shown in table 5 and Fig.

6.

Table 5. AUC,  $T_{max}$ ,  $C_{max}$ ,  $K_{el}$  and  $t_{1/2}$  on suppository composition

	AUC (hr-ng/ml)	MRT (hr)	$T_{max}$ (min.)	$C_{max}$ (ng/ml)	$K_{el}$ (hr <sup>-1</sup> )	$t_{1/2}$ (hr)
intravenous injection of propranolol solution	363.91 ±61.21	1.91 ±0.34	-	-	0.46 ±0.07	1.50 ±0.22
oral administration of propranolol solution	48.92 ±7.31	3.10 ±0.42	7.00 ±2.45	54.44 ±16.30	0.24 ±0.09	2.93 ±0.62
rectal administration of conventional suppository composition I	151.18 ±37.68	2.67 ±0.35	30.00 ±8.94	128.12 ±34.59	0.29 ±0.09	2.37 ±0.47
rectal administration of comparative suppository composition I	225.78 ±30.67	2.27 ±0.34	22.50 ±4.33	147.10 ±50.01	0.43 ±0.08	1.63 ±0.31
rectal administration of example 2	318.33 ±42.34	3.19 ±0.48	26.00 ±4.90	185.82 ±52.17	0.27 ±0.08	2.55 ±0.53

From the above results, it was revealed that the bioavailability on the suppository composition of Example 2 was significantly higher than that of comparative suppository composition and conventional

suppository composition, while having similar level of bioavailability when injected intravenously.

### CLAIMS

1. Suppository composition containing 0.1-10 weight part of a drug which undergoes the hepatic first-pass effect, 25-40 weight part of poloxamer and 0.1-1 weight part of hydrophilic natural polymers.

5

2. Suppository composition according to claim 1 wherein a drug which undergoes the hepatic first-pass effect is propranolol.

3. Suppository composition according to claim 1 wherein one or more poloxamers is/are selected from solid type, liquid type and paste type.


10

4. Suppository composition for rectal administration according to claim 1 wherein hydrophilic natural polymers are used from chitosan and sodium alginate independently or in a mixing form.

15

1 / 6


Fig. 1



- (A) F-127/F-68 (15/15 weight part)
- (B) F-127/F-68/PVP (15/15/1 weight part)
- (C) F-127/F-68/HPMC (15/15/1 weight part)
- (D) F-127/F-68/HPC (15/15/1 weight part)
- (E) F-127/F-68/sodium CMC (15/15/1 weight part)
- (F) F-127/F-68/carbopol (15/15/1 weight part)
- (G) F-127/F-68/polycarbophil (15/15/1 weight part)
- (H) F-127/F-68/sodium alginate (15/15/1 weight part)
- (I) F-127/F-68/chitosan (15/15/1 weight part)

2 / 6


Fig. 2



- (A) F-127/F-68 (15/15 weight part)
- (B) F-127/F-68/PVP (15/15/1 weight part)
- (C) F-127/F-68/HPMC (15/15/1 weight part)
- (D) F-127/F-68/HPC (15/15/1 weight part)
- (E) F-127/F-68/sodium CMC (15/15/1 weight part)
- (F) F-127/F-68/carbopol (15/15/1 weight part)
- (G) F-127/F-68/polycarbophil (15/15/1 weight part)
- (H) F-127/F-68/sodium alginate (15/15/1 weight part)
- (I) F-127/F-68/chitosan (15/15/1 weight part)

3 / 6


Fig. 3



- (A) F-127/F-68 (15/15 weight part)
- (B) F-127/F-68/PVP (15/15/1 weight part)
- (C) F-127/F-68/HPMC (15/15/1 weight part)
- (D) F-127/F-68/HPC (15/15/1 weight part)
- (E) F-127/F-68/sodium CMC (15/15/1 weight part)
- (F) F-127/F-68/carbopol (15/15/1 weight part%)
- (G) F-127/F-68/polycarbophil (15/15/1 weight part)
- (H) F-127/F-68/sodium alginate (15/15/1 weight part)
- (I) F-127/F-68/chitosan (15/15/1 weight part)

4 / 6

Fig. 4







- F-127/F-68/propranolol/sodium alginate (15/15/2/0.2 weight part)
- F-127/F-68/propranolol/sodium alginate (15/15/2/0.8 weight part)
- F-127/F-68/propranolol/polycarbophil (15/15/2/0.2 weight part)
- F-127/F-68/propranolol/polycarbophil (15/15/2/0.8 weight part)




5 / 6

Fig. 5



6 / 6

Fig. 6



- intravenous injection group of propranolol solution
- oral administration group of propranolol solution
- Δ- rectal administration group of conventional suppository composition I
- ▼- rectal administration group of the comparative suppository composition
- rectal administration group of the suppository composition of Example 2

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 97/00032

## A. CLASSIFICATION OF SUBJECT MATTER

IPC<sup>6</sup>: A 61 K 9/02, 47/34, 47/36

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC<sup>6</sup>: A 61 K 9/02, 47/34, 47/36

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

QUESTEL-WPIL

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 346 703 A (VIEGAS T.X. et al.) 13 September 1994 (13.09.94), claims 1-8,15-22,24,27,28; column 7, line 63 - column 8, line 10; column 8, lines 30-31; column 9, lines 4-21; column 10, lines 35-68.	1-4
X	US 5 292 516 A (VIEGAS T.X. et al.) 08 March 1994 (08.03.94), claims 1-4; column 8, line 55 - column 9, line 24; abstract; column 6, line 52 - column 7, line 68; example 3.	1-3
Y	EP 0 103 995 A2 (CILAG AG) 28 March 1984 (28.03.84), claims 1,2.	4
Y	US 4 946 870 A (PARTAIN E.M. et al.) 07 August 1990 (07.08.90), claims 1,11; column 8, lines 7-58.	4
X	WO 94/03 157 A1 (POLI INDUSTRIA CHIMICA S.P.A.) 17 February 1994 (17.02.94), claims 1,4,5,7-10,13; page 7, lines 9-14; page 8, lines 5-10.	1,3
A	WO 94/03 186 A1 (LABORATOIRES JEAN-PAUL MARTIN) 17 February 1994 (17.02.94), claims 1,4,6,14; page 3, lines 1-6; page 6, lines 1-18.	1,3,4

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 April 1997 (22.04.97)

Date of mailing of the international search report

13 May 1997 (13.05.97)

Name and mailing address of the ISA/ AT  
AUSTRIAN PATENT OFFICE  
Kohlmarkt 8-10  
A-1014 Vienna  
Facsimile No. 1/53424/535

Authorized officer  
Mazzucco  
Telephone No. 1/5337058/33

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR 97/00032

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EP 0 551 626 A1 (LEK) 21 July 1993 (21.07.93), abstract; claims 1-15; examples; (cited in the application).</p>	1-3

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.

PCT/KR 97/00032

la Recherchebericht angeführtes Patentedokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
US A 5346703	13-09-94	CA AA 2044878 DE CO 691220008 DE T2 691220008 EP A1 470703 EP A1 719545 EP B1 470703 JP A2 4230636 US A 5143731 US A 5277911 US A 5376693 US A 5124151 US A 5071644 US A 5077033 CA AA 2040460 DE CO 69124416 EP A1 455396 EP B1 455396 JP A2 4225914 US A 5298260 US A 533683 US A 5292516 US A 5306301 US A 5300295	08-02-92 17-10-96 30-01-97 12-02-92 03-07-96 11-09-96 19-08-93 01-09-92 11-01-94 27-12-94 10-12-91 01-12-91 02-11-91 13-03-97 06-11-91 29-01-97 14-08-93 29-03-94 14-01-97 08-03-94 26-04-94 05-04-94
US A 5292516	08-03-94	CA AA 2040460 DE CO 69124416 EP A1 455396 EP B1 455396 JP A2 4225914 CA AA 2044878 DE CO 691220008 DE T2 691220008 EP A1 470703 EP A1 719545 EP B1 470703 JP A2 4230636 US A 5077033 US A 5277911 US A 5376693 US A 5298260 US A 533683 US A 5306301 US A 5300295 US A 5124151 US A 5143731 US A 5346703 US A 5071644	02-11-91 13-03-97 06-11-91 29-01-97 14-08-93 08-03-93 17-10-96 30-01-97 12-02-92 03-07-96 11-09-96 19-08-93 01-12-91 11-01-94 27-12-94 29-03-94 14-01-97 26-04-94 05-04-94 23-06-93 01-09-96 13-09-94 10-12-91
EP A2 103995	28-03-84	AT E 51752 AU A1 18334/83 AU B2 557476 CA A1 1207231 DE CO 691220008 DK AO 6860/83 DK A 6860/83 DK B 162373 DK C 162373 EP A3 103995 EP B1 103995 FI A0 833019 FI B 833019 FI C 833019 GB A0 833019 GB A1 833019 GB B2 833019 HK A 833019 IL A 833019 IL A1 833019	15-04-80 01-03-84 24-12-86 09-07-86 17-05-90 23-08-83 23-02-84 23-11-84 09-03-90 19-09-95 11-04-90 23-08-93 23-11-91 10-03-92 20-09-94 21-03-94 24-12-86 08-07-86 01-08-90 13-11-88

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR 97/00032

US A	4946870	07-08-90	JP A2	59055817	31-03-84
			JP B4	60065330	26-01-94
			MY A	675787	31-12-87
			NO A	9330066	27-02-84
			NO B	1684005	11-11-91
			NO C	1684005	19-02-92
			PH A	18270	14-05-85
			SG A	766786	27-02-87
			ZA A	8306237	24-04-85
			US A	4698359	06-10-87
US A	4946870	07-08-90	AT E	144781	15-11-96
			CA A1	1283600	30-04-91
			DE C0	3751928	05-12-96
			DE T2	3751938	27-02-97
			EP A1	271551	22-06-88
			EP B1	271551	30-10-96
			JP T2	63503466	15-12-88
			JP B4	7076241	16-08-95
			WO A1	8707618	17-12-87
			US A	4929722	29-05-90
			US A	5300494	05-04-94
			AU A1	4449778	31-05-90
			AU B2	625075	02-07-92
			CA AA	2002404	08-05-90
			EP A2	368253	16-05-90
			EP A3	368253	29-09-91
			IL A0	922223	26-07-90
			IL A1	922223	08-07-93
			JP A2	2196738	03-08-90
			KR B1	9402657	28-03-94
WO A1	9403157	17-02-94	AT E	135566	15-04-96
			DE C0	69301920	25-04-96
			DE T2	69301920	08-08-96
			EP A1	652744	17-05-95
			EP B1	652744	20-03-96
			ES T3	2085166	16-05-96
			IT A0	92501831	28-07-92
			IT A	1255460	02-11-95
WO A1	9403186	17-02-94	EP A1	606457	20-07-94
			FR A1	2694194	04-02-94
			FR B1	2694194	04-11-94
			JP T2	8500578	23-01-96
EP A1	551626	21-07-93	keine - none - rien		